

A NEW GENERATION
OF PROTECTION

pure

HARD SURFACE

**DISINFECTANT, FUNGICIDE, VIRUCIDE* AND
FOOD CONTACT SURFACE SANITIZER**

IDEAL FOR CHILDCARE AND
EDUCATIONAL ENVIRONMENTS

**ELIMINATES MRSA,
INFLUENZA A (H1N1) VIRUS
AND NOROVIRUS**

ACTIVE INGREDIENTS

SILVER [†]	0.003%
CITRIC ACID	4.846%
OTHER INGREDIENTS	95.151%
TOTAL	100.000%

[†]Electrolytically generated silver ions stabilized
in citric acid as silver dihydrogen citrate

EPA REG. No. 72977-5-73912
EPA EST. No. 72977-CA-001

- KILLS COLD AND
FLU VIRUSES
- NO RINSE FORMULA
- ODORLESS

Net Contents **1 QT**
(32 FL OZ) 946 ML

KEEP OUT OF REACH OF CHILDREN

KILLS GERMS IN 30 SECONDS[‡]

DIRECTIONS FOR USE:

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

This product is a colorless, odorless, broad spectrum antimicrobial sanitizer, disinfectant and deodorizer. Proven to kill bacteria, fungus and viruses*, this product should be used on non-porous environmental hard surfaces in homes, schools, dorm rooms, daycare centers, nurseries, playgrounds, bathrooms, locker rooms, gyms, food storage areas, snack bars, supermarkets, kitchens, restaurants, lunchrooms and dining halls. Formulated to treat hard, non-porous environmental surfaces (painted, glazed tile, plastic, metal, glass) and objects including counters, desks, drinking fountains, kitchen surfaces, cutting boards, refrigerators, freezers, highchairs, children's toys, strollers, lunch boxes, coolers, cooking equipment, food trays, pet bowls, sinks and ice machines.

GENERAL INFORMATION: This product successfully killed the following organisms under AOAC protocols. In order to ensure that all organisms listed are killed, you must use the contact times as directed in the Application Instructions:

ORGANISM	KILL TIME
‡Pseudomonas aeruginosa	30 seconds
‡ Salmonella enterica	30 seconds
Staphylococcus aureus	2 minutes
Listeria monocytogenes	2 minutes
Vancomycin resistant Enterococcus faecium (VRE)	2 minutes
Methicillin resistant Staphylococcus aureus (MRSA)	2 minutes
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA)	2 minutes
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA-PVL)	2 minutes
Escherichia coli O157:H7	2 minutes
Acinetobacter baumannii	2 minutes
Campylobacter jejuni	2 minutes
Carbapenem resistant Escherichia coli	2 minutes
Carbapenem resistant Klebsiella pneumoniae	2 minutes
Carbapenem resistant Klebsiella pneumoniae, NDM-1 +	2 minutes
Trichophyton mentagrophytes (Athlete's Foot Fungus)	5 minutes
‡*HIV type 1	30 seconds
‡*Rotavirus	30 seconds
‡*Human Coronavirus	30 seconds
‡*Influenza A (H1N1) • ‡*Swine Influenza A (H1N1)	30 seconds
‡*Respiratory Syncytial Virus	30 seconds
‡*Adenovirus Type 2	30 seconds
‡*Influenza A • ‡*Avian Influenza A	30 seconds
*Hepatitis B Virus (HBV)	60 seconds
*Hepatitis C Virus (HCV)	60 seconds
*Norovirus • *Murine Norovirus	60 seconds
*Herpes Simplex Type 1	60 seconds
*Rhinovirus	60 seconds
*Polio Type 2	60 seconds

Distributed by
PURE BIOSCIENCE
1725 Gillespie Way
El Cajon, CA 92020



MADE IN THE USA



PEEL HERE

APPLICATION INSTRUCTIONS:

Pre-clean surfaces prior to using this product. You may use this product for pre-cleaning.

GENERAL CLEANING: Apply to surface until thoroughly wet then wipe the surface clean.

GENERAL DISINFECTION: To kill bacteria, apply this product to the surface until thoroughly wet for 2 minutes. The surface may then be wiped dry with a clean towel. When used as directed, this product provides residual protection from *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella enterica* up to 24 hours after initial application. Do not touch treated surface after application if residual protection is to be maintained.

FUNGUS CONTROL: To kill fungus, apply this product to the surface until thoroughly wet for 5 minutes. The surface may then be wiped dry with a clean towel. Re-apply when cleaning or when new growth appears.

***VIRAL CONTROL:** To kill viruses, apply this product to the surface until thoroughly wet for 1 minute. The surface may then be wiped dry with a clean towel.

This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

Kills HIV-1, HBV and HCV on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings (or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV, HBV or HCV); **Instructions for Cleaning and Decontamination Against HIV, HBV and HCV** on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids: **Personal Protection:** When handling items soiled with blood or body fluids, use appropriate barrier protection such as latex gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant. **Contact Time:** Apply to area to be treated. Allow the surface to remain wet for 30 seconds to kill HIV-1. Use 1 minute for HBV and HCV. The surface may then be wiped dry with a clean towel. These contact times will not control all organisms listed on this label. Refer to application instructions for other organisms. **Disposal of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

U.S. Patents: 6,197,814; 6,583,176; 7,261,905; 7,763,297; 7,803,407

Other patents pending

**APPLICATION
INSTRUCTIONS (CONT'D):**

SANITIZATION OF FOOD CONTACT SURFACES:

Do not use this product on utensils, dishes or glassware.

Spray, pour or spread this product on surface until thoroughly wet.
Let stand for 60 seconds and wipe with a clean towel or allow to air dry.
No rinsing is required.

Kills 99.999% of *Escherichia coli* and *Staphylococcus aureus*.

TO REFILL SPRAY BOTTLES:

1. Remove trigger sprayer from empty bottle.
2. Remove cap from refill and pour contents directly into empty bottle.
3. Replace trigger sprayer and use as you normally would.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area away from direct sunlight at temperatures above freezing.

PESTICIDE DISPOSAL: Nonrefillable Container. Refill only with this product. Do not reuse or refill except as described in the directions for use. If empty: Place in trash or offer for recycling if available. Rinse thoroughly before discarding in trash or recycling.

IN CASE OF EMERGENCY: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC 1-800-424-9300 for emergency medical treatment information.